

YOUR GLBT GUIDE TO

Key West



Q MAGAZINE

Volume 8 - Issue 5

May 2013



ASK NANA



THE INQUISITOR



MAN ABOUT TOWN



DANCE

THE AFTER "T" EVENT

EVERY SUNDAY 6-10PM

MUSIC BY **DJ NEIL**

JESSICA DEVERAUX VARIETY HOUR AT 8PM

SEXY HOT DANCERS

724 DUVAL • KEY WEST

Bourbon St. Pub

BourbonStPub.com





Q MAGAZINE



Welcome to
Paradise!

This month we debut two new features in Q Magazine.

First is Zachary Moses with Man About Town. This monthly trip through Key West will bring you to the best places to eat, drink and shop!

We also have The InQuisitor. This column will dish all the hottest gossip going on around the community!

Enjoy!

Contributors:

Neil Chamberlain
Zachary Moses
Brad Loekle
Nana

For advertising information,
please see us online at:
www.QKeyWest.com

Published Monthly by:
The Gay Rag, Inc.
Key West, FL 33040

info@keywestgayrag.com

305.396.8227

Visit us online at www.QKeyWest.com

What is STRIBILD?

STRIBILD is a prescription medicine used to treat HIV-1 in adults who have never taken HIV-1 medicines before. It combines 4 medicines into 1 pill to be taken once a day with food. STRIBILD is a complete single-tablet regimen and should not be used with other HIV-1 medicines.

STRIBILD does not cure HIV-1 infection or AIDS. To control HIV-1 infection and decrease HIV-related illnesses you must keep taking STRIBILD. Ask your healthcare provider if you have questions about how to reduce the risk of passing HIV-1 to others. Always practice safer sex and use condoms to lower the chance of sexual contact with body fluids. Never reuse or share needles or other items that have body fluids on them.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about STRIBILD?

STRIBILD can cause serious side effects:

- **Build-up of an acid in your blood (lactic acidosis)**, which is a serious medical emergency. Symptoms of lactic acidosis include feeling very weak or tired, unusual (not normal) muscle pain, trouble breathing, stomach pain with nausea or vomiting, feeling cold especially in your arms and legs, feeling dizzy or lightheaded, and/or a fast or irregular heartbeat.
- **Serious liver problems.** The liver may become large (hepatomegaly) and fatty (steatosis). Symptoms of liver problems include your skin or the white part of your eyes turns yellow (jaundice), dark "tea-colored" urine, light-colored bowel movements (stools), loss of appetite for several days or longer, nausea, and/or stomach pain.
- **You may be more likely to get lactic acidosis or serious liver problems** if you are female, very overweight (obese), or have been taking STRIBILD for a long time. In some cases, these serious conditions have led to death. Call your healthcare provider right away if you have any symptoms of these conditions.

- **Worsening of hepatitis B (HBV) infection.** If you also have HBV and stop taking STRIBILD, your hepatitis may suddenly get worse. Do not stop taking STRIBILD without first talking to your healthcare provider, as they will need to monitor your health. STRIBILD is not approved for the treatment of HBV.

Who should not take STRIBILD?

Do not take STRIBILD if you:

- **Take a medicine that contains:** afuzosin, dihydroergotamine, ergotamine, methylergonovine, cisapride, lovastatin, simvastatin, pimozide, sildenafil when used for lung problems (Revatio®), triazolam, oral midazolam, rifampin or the herb St. John's wort.
- **For a list of brand names for these medicines,** please see the Brief Summary on the following pages.
- **Take any other medicines to treat HIV-1 infection,** or the medicine *adefovir* (Hepsera®).

What are the other possible side effects of STRIBILD?

Serious side effects of STRIBILD may also include:

- **New or worse kidney problems, including kidney failure.** Your healthcare provider should do regular blood and urine tests to check your kidneys before and during treatment with STRIBILD. If you develop kidney problems, your healthcare provider may tell you to stop taking STRIBILD.
- **Bone problems,** including bone pain or bones getting soft or thin, which may lead to fractures. Your healthcare provider may do tests to check your bones.
- **Changes in body fat** can happen in people taking HIV-1 medicines.
- **Changes in your immune system.** Your immune system may get stronger and begin to fight infections. Tell your healthcare provider if you have any new symptoms after you start taking STRIBILD.

The most common side effects of STRIBILD include nausea and diarrhea. Tell your healthcare provider if you have any side effects that bother you or don't go away.

What should I tell my healthcare provider before taking STRIBILD?

- **All your health problems.** Be sure to tell your healthcare provider if you have or had any kidney, bone, or liver problems, including hepatitis virus infection.
 - **All the medicines you take,** including prescription and nonprescription medicines, vitamins, and herbal supplements. STRIBILD may affect the way other medicines work, and other medicines may affect how STRIBILD works. Keep a list of all your medicines and show it to your healthcare provider and pharmacist. Do not start any new medicines while taking STRIBILD without first talking with your healthcare provider.
 - **If you take hormone-based birth control** (pills, patches, rings, shots, etc.).
 - **If you take antacids.** Take antacids at least 2 hours before or after you take STRIBILD.
 - **If you are pregnant or plan to become pregnant.** It is not known if STRIBILD can harm your unborn baby. Tell your healthcare provider if you become pregnant while taking STRIBILD.
 - **If you are breastfeeding** (nursing) or plan to breastfeed. Do not breastfeed. HIV-1 can be passed to the baby in breast milk. Also, some medicines in STRIBILD can pass into breast milk, and it is not known if this can harm the baby.
- You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.**

Please see Brief Summary of full Prescribing Information with important warnings on the following pages.



STRIBILD is a prescription medicine used as a complete single-tablet regimen to treat HIV-1 in adults who have never taken HIV-1 medicines before. STRIBILD does not cure HIV-1 or AIDS.

I started my personal revolution

Talk to your healthcare provider about starting treatment.

STRIBILD is a complete HIV-1 treatment in **1 pill**, once a day.

Ask if it's right for you.

STRIBILD™ 

elvitegravir 150mg/ cobicistat 150mg/ emtricitabine
200mg/ tenofovir disoproxil fumarate 300mg tablets

Patient Information

STRIBILD™ (STRY-bild)

(elvitegravir 150 mg/cobicistat 150 mg/
emtricitabine 200 mg/tenofovir disoproxil
fumarate 300 mg) tablets

Brief summary of full Prescribing Information.
For more information, please see the full Prescribing
Information, including Patient Information.

What is STRIBILD?

- **STRIBILD is a prescription medicine used to treat HIV-1 in adults who have never taken HIV-1 medicines before.** STRIBILD is a complete regimen and should not be used with other HIV-1 medicines.
- **STRIBILD does not cure HIV-1 or AIDS.** You must stay on continuous HIV-1 therapy to control HIV-1 infection and decrease HIV-related illnesses.
- **Ask your healthcare provider about how to prevent passing HIV-1 to others.** Do not share or reuse needles, injection equipment, or personal items that can have blood or body fluids on them. Do not have sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.

What is the most important information I should know about STRIBILD?

STRIBILD can cause serious side effects, including:

1. **Build-up of lactic acid in your blood (lactic acidosis).** Lactic acidosis can happen in some people who take STRIBILD or similar (nucleoside analogs) medicines. Lactic acidosis is a serious medical emergency that can lead to death. Lactic acidosis can be hard to identify early, because the symptoms could seem like symptoms of other health problems. **Call your healthcare provider right away if you get any of the following symptoms which could be signs of lactic acidosis:**

- feel very weak or tired
- have unusual (not normal) muscle pain
- have trouble breathing
- have stomach pain with nausea or vomiting
- feel cold, especially in your arms and legs

- feel dizzy or lightheaded
 - have a fast or irregular heartbeat
2. **Severe liver problems.** Severe liver problems can happen in people who take STRIBILD. In some cases, these liver problems can lead to death. Your liver may become large (hepatomegaly) and you may develop fat in your liver (steatosis). **Call your healthcare provider right away if you get any of the following symptoms of liver problems:**

- your skin or the white part of your eyes turns yellow (jaundice)
- dark “tea-colored” urine
- light-colored bowel movements (stools)
- loss of appetite for several days or longer
- nausea
- stomach pain

You may be more likely to get lactic acidosis or severe liver problems if you are female, very overweight (obese), or have been taking STRIBILD for a long time.

3. **Worsening of Hepatitis B infection.** If you have hepatitis B virus (HBV) infection and take STRIBILD, your HBV may get worse (flare-up) if you stop taking STRIBILD. A “flare-up” is when your HBV infection suddenly returns in a worse way than before.

- Do not run out of STRIBILD. Refill your prescription or talk to your healthcare provider before your STRIBILD is all gone
- Do not stop taking STRIBILD without first talking to your healthcare provider
- If you stop taking STRIBILD, your healthcare provider will need to check your health often and do blood tests regularly for several months to check your HBV infection. Tell your healthcare provider about any new or unusual symptoms you may have after you stop taking STRIBILD

Who should not take STRIBILD?

Do not take STRIBILD if you also take a medicine that contains:

- adefovir (Hepsera®)
- alfuzosin hydrochloride (Uroxatral®)
- cisapride (Propulsid®, Propulsid Quicksolv®)
- ergot-containing medicines, including:

dihydroergotamine mesylate (D.H.E. 45[®], Migranal[®]), ergotamine tartrate (Cafergot[®], Migergot[®], Ergostat[®], Medihaler Ergotamine[®], Wigraine[®], Wigrettes[®]), and methylergonovine maleate (Ergotrate[®], Methergine[®])

- lovastatin (Advicor[®], Altoprev[®], Mevacor[®])
- oral midazolam
- pimozide (Orap[®])
- rifampin (Rifadin[®], Rifamate[®], Rifater[®], Rimactane[®])
- sildenafil (Revatio[®]), when used for treating lung problems
- simvastatin (Simcor[®], Vytorin[®], Zocor[®])
- triazolam (Halcion[®])
- the herb St. John's wort

Do not take STRIBILD if you also take any other HIV-1 medicines, including:

- Other medicines that contain tenofovir (Atripla[®], Complera[®], Viread[®], Truvada[®])
- Other medicines that contain emtricitabine, lamivudine, or ritonavir (Combivir[®], Emtriva[®], Epivir[®] or Epivir-HBV[®], Epzicom[®], Kaletra[®], Norvir[®], Trizivir[®])

STRIBILD is not for use in people who are less than 18 years old.

What are the possible side effects of STRIBILD?

STRIBILD may cause the following serious side effects:

- **See "What is the most important information I should know about STRIBILD?"**
- **New or worse kidney problems, including kidney failure.** Your healthcare provider should do blood and urine tests to check your kidneys before you start and while you are taking STRIBILD. Your healthcare provider may tell you to stop taking STRIBILD if you develop new or worse kidney problems.
- **Bone problems** can happen in some people who take STRIBILD. Bone problems include bone pain, softening or thinning (which may lead to fractures). Your healthcare provider may need to do tests to check your bones.
- **Changes in body fat** can happen in people who take HIV-1 medicine. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the middle of your body (trunk). Loss of fat from the legs, arms and face may also

happen. The exact cause and long-term health effects of these conditions are not known.

- **Changes in your immune system** (Immune Reconstitution Syndrome) can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having any new symptoms after starting your HIV-1 medicine.

The most common side effects of STRIBILD include:

- Nausea
- Diarrhea

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

- These are not all the possible side effects of STRIBILD. For more information, ask your healthcare provider.
- Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What should I tell my healthcare provider before taking STRIBILD?

Tell your healthcare provider about all your medical conditions, including:

- If you have or had any kidney, bone, or liver problems, including hepatitis B infection
- If you are pregnant or plan to become pregnant. It is not known if STRIBILD can harm your unborn baby. Tell your healthcare provider if you become pregnant while taking STRIBILD.
 - There is a pregnancy registry for women who take antiviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk with your healthcare provider about how you can take part in this registry.
- If you are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed if you take STRIBILD.
 - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.
 - Two of the medicines in STRIBILD can pass to your baby in your breast milk. It is not known if the other medicines in STRIBILD can pass into your breast milk.

- Talk with your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements:

- STRIBILD may affect the way other medicines work, and other medicines may affect how STRIBILD works.

- Be sure to tell your healthcare provider if you take any of the following medicines:

- Hormone-based birth control (pills, patches, rings, shots, etc)
- Antacid medicines that contains aluminum, magnesium hydroxide, or calcium carbonate. Take antacids at least 2 hours before or after you take STRIBILD
- Medicines to treat depression, organ transplant rejection, or high blood pressure
- amiodarone (Cordarone[®], Pacerone[®])
- atorvastatin (Lipitor[®], Caduet[®])
- bepridil hydrochloric (Vascor[®], Bepadin[®])
- bosentan (Tracleer[®])
- buspirone
- carbamazepine (Carbatrol[®], Epitol[®], Equetro[®], Tegreto[®])
- clarithromycin (Biaxin[®], Prevpac[®])
- clonazepam (Klonopin[®])
- clorazepate (Gen-xene[®], Tranxene[®])
- colchicine (Colcrys[®])
- medicines that contain dexamethasone
- diazepam (Valium[®])
- digoxin (Lanoxin[®])
- disopyramide (Norpace[®])
- estazolam
- ethosuximide (Zarontin[®])
- flecainide (Tambocor[®])
- flurazepam
- fluticasone (Flovent[®], Flonase[®], Flovent[®] Diskus, Flovent[®] HFA, Veramyst[®])
- itraconazole (Sporanox[®])
- ketoconazole (Nizoral[®])
- lidocaine (Xylocaine[®])
- mexiletine
- oxcarbazepine (Trileptal[®])

- perphenazine
- phenobarbital (Luminal[®])
- phenytoin (Dilantin[®], Phenytek[®])
- propafenone (Rythmol[®])
- quinidine (Neudexta[®])
- rifabutin (Mycobutin[®])
- rifapentine (Priftin[®])
- risperidone (Risperdal[®], Risperdal Consta[®])
- salmeterol (Serevent[®]) or salmeterol when taken in combination with fluticasone (Advair Diskus[®], Advair HFA[®])
- sildenafil (Viagra[®]), tadalafil (Cialis[®]) or vardenafil (Levitra[®], Staxyn[®]), for the treatment of erectile dysfunction (ED). If you get dizzy or faint (low blood pressure), have vision changes or have an erection that last longer than 4 hours, call your healthcare provider or get medical help right away.
- tadalafil (Adcirca[®]), for the treatment of pulmonary arterial hypertension
- telithromycin (Ketek[®])
- thioridazine
- voriconazole (Vfend[®])
- warfarin (Coumadin[®], Jantoven[®])
- zolpidem (Ambien[®], Edlular[®], Intermezzo[®], Zolpimist[®])

Know the medicines you take. Keep a list of all your medicines and show it to your healthcare provider and pharmacist when you get a new medicine. Do not start any new medicines while you are taking STRIBILD without first talking with your healthcare provider.

Keep STRIBILD and all medicines out of reach of children.

This Brief Summary summarizes the most important information about STRIBILD. If you would like more information, talk with your healthcare provider. You can also ask your healthcare provider or pharmacist for information about STRIBILD that is written for health professionals, or call 1-800-445-3235 or go to www.STRIBILD.com.

Issued: August 2012



COMPLERA, EMTRIVA, GILEAD, the GILEAD Logo, GS1, HEPSERA, STRIBILD, the STRIBILD Logo, TRUVADA, and VIREAD are trademarks of Gilead Sciences, Inc. or its related companies. ATRIPLA is a trademark of Bristol-Myers Squibb & Gilead Sciences, LLC. All other marks referenced herein are the property of their respective owners.

SNAP



SNAP



Photos By
Larry
Blackburn



WEEKLY EVENTS



MONDAY

801

Drag Shows at 9 PM and 11 PM

Bobby's

Karaoke starting at 9:30 PM

Bourbon

Dancers on the bar starting at 10 PM

TUESDAY

801

Drag Shows at 9 PM and 11 PM

Bourbon

Dancers on the bar starting at 10 PM



801- 801 Duval St.



Bobby's- 900 Simonton St.



Bourbon- 724 Duval St.

W E D N E S D A Y

801

Drag Shows at 9 PM and 11 PM

Bourbon

Dancers on the bar starting at 10 PM

The Little Red Store

25% off from 6 PM - 8 PM

T H U R S D A Y

801

Karaoke starting at 5 PM

Drag Shows at 9 PM and 11 PM

Bobby's

Karaoke starting at 9:30 PM

Bourbon

Dancers on the bar starting at 10 PM

F R I D A Y

801

*Drag Shows at 5 PM, 9 PM
and 11 PM*

Bobby's

Karaoke starting at 9:30 PM

Bourbon

Dancers on the bar starting at 10 PM

Stripper Battle starting at Midnight

One Saloon

*Cock Shock contest starting at
Midnight*



Island House- 1129 Flemingl St.

S A T U R D A Y

801

*Drag Shows at 5PM, 9 PM
and 11-PM*

Bourbon

Pool Party starting at 12 PM

Dancers on the bar starting at 10 PM

Amateur Strip at Midnight

S U N D A Y

801

Karaoke starting at 5 PM

Gay Bingo starting at 5 PM

Drag Shows at 9 PM and 11 PM

Bobby's

Karaoke starting at 9:30 PM

Bourbon

After Tea Dance starting at 6PM

Dancers on the bar starting at 10 PM

Island House

Pre-Tea Pool Party starting at Noon



One Saloon- 514 Petronia St.

801 CABARET

GIRLS GIRLS GIRLS

Key West



NIGHTLY 9 & 11 PM

801 DUVAL • KEY WEST

801 Bourbon Bar

801 Bourbon Bar



801 Bourbon PRESENTS

MARTINI WEDNESDAYS

\$5 MARTINIS | EACH AND EVERY WEDNESDAY NIGHT AT THE MAIN BAR

21+ MUST HAVE ID TIME 10PM-2AM

LOCATION 801 DUVAL ST., KEY WEST, 33040

WWW.801BOURBON.COM | 305.731.9055

GASSY & JEFF

DRAG-KARAOKE SUNDAYS

THE SINGING BEGINS @ 4PM

\$5 LONG ISLAND ICED TEAS ALL DAY LONG!

801 BOURBON
801 DUVAL STREET
KEY WEST, FL 33040
CALL: 305.731.9055
WWW.801BOURBON.COM

WEDNESDAYS 2012 6PM

KARAOKE NIGHTS

KARAOKE BEGINS AT 6PM MARTINI SPECIALS BEGINS AT 10PM

Emcee Jeff 6pm-8pm

Emcee Rhett 8pm-10pm

801 Bourbon Bar

COME SING WITH THE BEST | WITH OVER 50,000 SONGS TO CHOOSE FROM
COME FOR THE KARAOKE... STAY FOR THE MARTINI'S

801 DUVAL STREET KEY WEST, FL | WWW.801BOURBON.COM | PH: 305.731.9055

BINGO

EVERY SUNDAY

QNTCH

BINGO
28 54 12 59 43

BINGO
4 18 37 59

21 5 18

ONE OF KEY WEST'S LONGEST RUNNING BENEFIT
DOORS OPEN AT 4PM - GAMES START AT 6PM

801 CABARE LOUNGE | 801 DUVAL STREET | WWW.801BOURBON.COM



The one for me

Patient model. Pill shown is not actual size.

What is COMPLERA?

COMPLERA® is a prescription HIV medicine that is used as a complete regimen to treat HIV-1 in adults who have **never** taken HIV medicines before **and** who have an amount of HIV in their blood (this is called “viral load”) that is no more than 100,000 copies/mL. COMPLERA contains 3 medicines — rilpivirine, emtricitabine and tenofovir disoproxil fumarate. It is not known if COMPLERA is safe and effective in children under the age of 18 years.

COMPLERA® does not cure HIV-1 infection or AIDS. To control HIV-1 infection and decrease HIV-related illnesses you must keep taking COMPLERA. Avoid doing things that can spread HIV-1 to others: always practice safer sex and use condoms to lower the chance of sexual contact with body fluids; never reuse or share needles or other items that have body fluids on them, do not share personal items that may contain bodily fluids. Ask your healthcare provider if you have questions about how to reduce the risk of passing HIV-1 to others.

IMPORTANT SAFETY INFORMATION

What is the most important information you should know about COMPLERA?

COMPLERA® can cause serious side effects:

- **Build-up of an acid in your blood (lactic acidosis)**, which is a serious medical emergency. Symptoms of lactic acidosis include feeling very weak or tired, unusual (not normal) muscle pain, trouble breathing, stomach pain with nausea or vomiting, feeling cold, especially in your arms and legs, feeling dizzy or lightheaded, and/or a fast or irregular heartbeat.
- **Serious liver problems.** The liver may become large (hepatomegaly) and fatty (steatosis). Symptoms of liver problems include your skin or the white part of your eyes turns yellow (jaundice), dark “tea-colored” urine, light-colored bowel movements (stools), loss of appetite for several days or longer, nausea, and/or stomach pain.
- **You may be more likely to get lactic acidosis or serious liver problems** if you are female, very overweight (obese), or have been taking COMPLERA for a long time. *In some cases, these serious conditions have led to death.* Call your healthcare provider right away if you have any symptoms of these conditions.

- **Worsening of hepatitis B (HBV) infection.** If you also have HBV and stop taking COMPLERA, your hepatitis may suddenly get worse. Do not stop taking COMPLERA without first talking to your healthcare provider, as they will need to monitor your health. COMPLERA is not approved for the treatment of HBV.

Who should not take COMPLERA?

Do not take COMPLERA if you have ever taken other anti-HIV medicines. COMPLERA may change the effect of other medicines and may cause serious side effects. Your healthcare provider may change your other medicines or change their doses. **Do not take COMPLERA if you also take these medicines:**

- **anti-seizure medicines:** carbamazepine (Carbatrol, Equetro, Tegretol, Tegretol-XR, Teril, Eptol); oxcarbazepine (Trileptal), phenobarbital (Luminal), phenytoin (Dilantin, Dilantin-125, Phenytek)
- **anti-tuberculosis medicines:** rifabutin (Mycobutin), rifampin (Rifater, Rifamate, Rimactane, Rifadin) and rifapentine (Priftin)
- **proton pump inhibitors for stomach or intestinal problems:** esomeprazole (Nexium, Vimovo), lansoprazole (Prevacid), dexlansoprazole (Dexilant), omeprazole (Prilosec), pantoprazole sodium (Protonix), rabeprazole (Aciphex)
- **more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate**
- **St. John’s wort (*Hypericum perforatum*)**

If you are taking COMPLERA you should not take other HIV medicines or other medicines containing tenofovir (Viread, Truvada, Stribild or Atripla); other medicines containing emtricitabine or lamivudine (Emtriva, Combivir, Epivir, Epivir-HBV, Epcicom, Trizivir, Atripla, Stribild or Truvada); rilpivirine (Edurant) or adefovir (Hepsera).

In addition, tell your healthcare provider if you are taking the following medications because they may interfere with how COMPLERA works and may cause side effects:

- certain antacid medicines containing aluminum, magnesium hydroxide, or calcium carbonate (examples: Rolaids, TUMS). **These medicines must be taken at least 2 hours before or 4 hours after COMPLERA.**
- medicines to block stomach acid including cimetidine (Tagamet), famotidine (Pepcid), nizatidine (Axid), or ranitidine HCL (Zantac). **These medicines must be taken at least 12 hours before or 4 hours after COMPLERA.**

COMPLERA. A complete HIV treatment in only 1 pill a day.

COMPLERA is for adults who have never taken HIV-1 medicines before and have no more than 100,000 copies/mL of virus in their blood.

Ask your healthcare provider if it's the one for you.

- any of these medicines: clarithromycin (Biaxin); erythromycin (E-Mycin, Eryc, Ery-Tab, PCE, Pediazole, Ilosone), fluconazole (Diflucan), itraconazole (Sporanox), ketoconazole (Nizoral) methadone (Dolophine); posaconazole (Noxafil), telithromycin (Ketek) or voriconazole (Vfend).
- medicines that are eliminated by the kidneys like acyclovir (Zovirax), cidofovir (Vistide), ganciclovir (Cytovene IV, Vitrasert), valacyclovir (Valtrex) and valganciclovir (Valcyte).

These are not all the medicines that may cause problems if you take COMPLERA. Tell your healthcare provider about all prescription and nonprescription medicines, vitamins, or herbal supplements you are taking or plan to take.

Before taking COMPLERA, tell your healthcare provider if you:

- Have **liver problems**, including hepatitis B or C virus infection, or have abnormal liver tests
- Have **kidney problems**
- Have ever had a **mental health problem**
- Have **bone problems**
- Are **pregnant or planning to become pregnant**. It is not known if COMPLERA can harm your unborn child
- Are **breastfeeding**: Women with HIV should not breastfeed because they can pass HIV through their milk to the baby. Also, COMPLERA may pass through breast milk and could cause harm to the baby

COMPLERA can cause additional serious side effects:

- **New or worsening kidney problems, including kidney failure.** If you have had kidney problems, or take other medicines that may cause kidney problems, your healthcare provider may need to do regular blood tests.
- **Depression or mood changes.** Tell your healthcare provider right away if you have any of the following symptoms: feeling sad or hopeless, feeling anxious or restless, have thoughts of hurting yourself (suicide) or have tried to hurt yourself.
- **Changes in liver enzymes:** People who have had hepatitis B or C, or who have had changes in their liver function tests in the past may have an increased risk for liver problems while taking COMPLERA. Some people without prior liver disease may also be at risk. Your healthcare provider may need to check your liver enzymes before and during treatment with COMPLERA.

- **Bone problems** can happen in some people who take COMPLERA. Bone problems include bone pain, softening or thinning (which may lead to fractures). Your healthcare provider may need to do additional tests to check your bones.
- **Changes in body fat** can happen in people taking HIV medicine.
- **Changes in your immune system.** Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting COMPLERA.

The most common side effects reported with COMPLERA are trouble sleeping (insomnia), abnormal dreams, headache, dizziness, diarrhea, nausea, rash, tiredness, and depression. Some side effects also reported include vomiting, stomach pain or discomfort, skin discoloration (small spots or freckles) and pain.

This is not a complete list of side effects. Tell your healthcare provider or pharmacist if you notice any side effects while taking COMPLERA, and call your healthcare provider for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <http://www.fda.gov/medwatch> or call 1-800-FDA-1088.

Additional Information about taking COMPLERA:

- **Always take COMPLERA exactly as your healthcare provider tells you to take it.**
- **Take COMPLERA with a meal.** Taking COMPLERA with a meal is important to help get the right amount of medicine in your body. (A protein drink does not replace a meal).

Stay under the care of your healthcare provider during treatment with COMPLERA and see your healthcare provider regularly.

Please see Brief Summary of full Prescribing Information with important warnings on the following pages.



COMPLERA[®]
emtricitabine 200mg/rilpivirine 25mg/
tenofovir disoproxil fumarate 300mg tablets

Learn more at www.COMPLERA.com

Brief Summary of full Prescribing Information

COMPLERA® (kom-PLEH-rah)

(emtricitabine, rilpivirine, tenofovir disoproxil fumarate) tablets

Brief summary of full Prescribing Information. For more information, please see the full Prescribing Information including Patient Information.

What is COMPLERA?

- **COMPLERA** is a prescription HIV (Human Immunodeficiency Virus) medicine that is used to treat HIV-1 in adults
 - who have **never** taken HIV medicines before, and
 - who have an amount of HIV in their blood (this is called ‘viral load’) that is no more than 100,000 copies/mL. Your healthcare provider will measure your viral load.

(HIV is the virus that causes AIDS (Acquired Immunodeficiency Syndrome)).

- COMPLERA contains 3 medicines – rilpivirine, emtricitabine, tenofovir disoproxil fumarate – combined in one tablet. It is a complete regimen to treat HIV-1 infection and should not be used with other HIV medicines.
- It is not known if COMPLERA is safe and effective in children under the age of 18 years old.
- **COMPLERA does not cure HIV infection or AIDS.** You must stay on continuous therapy to control HIV infection and decrease HIV-related illnesses.
- **Ask your healthcare provider if you have any questions about how to prevent passing HIV to other people.** Do not share or re-use needles or other injection equipment, and do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal fluids or blood.

What is the most important information I should know about COMPLERA?

COMPLERA can cause serious side effects, including:

- **Build-up of lactic acid in your blood (lactic acidosis).** Lactic acidosis can happen in some people who take COMPLERA or similar (nucleoside analogs) medicines. Lactic acidosis is a serious medical emergency that can lead to death. Lactic acidosis can be hard to identify early, because the symptoms could seem like symptoms of other health problems. **Call your healthcare provider right away if you get any of the following symptoms which could be signs of lactic acidosis:**
 - feel very weak or tired
 - have unusual (not normal) muscle pain
 - have trouble breathing

- have stomach pain with nausea (feeling sick to your stomach) or vomiting
- feel cold, especially in your arms and legs
- feel dizzy or lightheaded
- have a fast or irregular heartbeat

- **Severe liver problems.** Severe liver problems can happen in people who take COMPLERA. In some cases, these liver problems can lead to death. Your liver may become large (hepatomegaly) and you may develop fat in your liver (steatosis). **Call your healthcare provider right away if you get any of the following symptoms of liver problems:**

- your skin or the white part of your eyes turns yellow (jaundice)
- dark “tea-colored” urine
- light-colored bowel movements (stools)
- loss of appetite for several days or longer
- nausea
- stomach pain

- **You may be more likely to get lactic acidosis or severe liver problems if you are female, very overweight (obese), or have been taking COMPLERA for a long time.**

- **Worsening of Hepatitis B infection.** If you have hepatitis B virus (HBV) infection and take COMPLERA, your HBV may get worse (flare-up) if you stop taking COMPLERA. A “flare-up” is when your HBV infection suddenly returns in a worse way than before. COMPLERA is not approved for the treatment of HBV, so you must discuss your HBV with your healthcare provider.

- Do not let your COMPLERA run out. Refill your prescription or talk to your healthcare provider before your COMPLERA is all gone.
- Do not stop taking COMPLERA without first talking to your healthcare provider.
- If you stop taking COMPLERA, your healthcare provider will need to check your health often and do blood tests regularly to check your HBV infection. Tell your healthcare provider about any new or unusual symptoms you may have after you stop taking COMPLERA.

Who should not take COMPLERA?

Do not take COMPLERA if:

- your HIV infection has been previously treated with HIV medicines.
- you are taking any of the following medicines:
 - **anti-seizure medicines:** carbamazepine (Carbatrol, Equetro, Tegretol, Tegretol-XR, Teril, Eptol); oxcarbazepine (Trileptal); phenobarbital (Luminal); phenytoin (Dilantin, Dilantin-125, Phenytek)

- **anti-tuberculosis (anti-TB) medicines:** rifabutin (Mycobutin); rifampin (Rifater, Rifamate, Rimactane, Rifadin); rifapentine (Priftin)
- **proton pump inhibitor (PPI) medicine** for certain stomach or intestinal problems: esomeprazole (Nexium, Vimovo); lansoprazole (Prevacid); dexlansoprazole (Dexilant); omeprazole (Prilosec, Zegerid); pantoprazole sodium (Protonix); rabeprazole (Aciphex)
- more than 1 dose of the steroid medicine dexamethasone or dexamethasone sodium phosphate
- St. John's wort (*Hypericum perforatum*)

• If you take COMPLERA, you should not take:

- Other medicines that contain tenofovir (Atripla, Stribild, Truvada, Viread)
- Other medicines that contain emtricitabine or lamivudine (Combivir, Emtriva, Eпивir or Eпивir-HBV, Epicom, Trizivir, Atripla, Truvada, Stribild)
- rilpivirine (Edurant)
- adefovir (Hepsera)

What should I tell my healthcare provider before taking COMPLERA?

Before you take COMPLERA, tell your healthcare provider if you:

- have or had liver problems, including hepatitis B or C virus infection, kidney problems, mental health problem or bone problems
- are pregnant or plan to become pregnant. It is not known if COMPLERA can harm your unborn child.

Pregnancy Registry. There is a pregnancy registry for women who take antiviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.

- are breast-feeding or plan to breast-feed. **You should not breastfeed if you have HIV because of the risk of passing HIV to your baby.** Do not breastfeed if you are taking COMPLERA. At least two of the medicines contained in COMPLERA can be passed to your baby in your breast milk. We do not know whether this could harm your baby. Talk to your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements.

- **COMPLERA may affect the way other medicines work, and other medicines may affect how COMPLERA works, and may cause serious side effects.** If you take certain medicines with COMPLERA, the amount of COMPLERA in your body may be too low and it may not work to help control your HIV infection. The HIV virus in your body may become resistant to COMPLERA or other HIV medicines that are like it.

Especially tell your healthcare provider if you take:

- an antacid medicine that contains aluminum, magnesium hydroxide, or calcium carbonate. If you take an antacid during treatment with COMPLERA, take the antacid **at least 2 hours before or at least 4 hours after** you take COMPLERA.
- a medicine to block the acid in your stomach, including cimetidine (Tagamet), famotidine (Pepcid), nizatidine (Axid), or ranitidine hydrochloride (Zantac). If you take one of these medicines during treatment with COMPLERA, take the acid blocker **at least 12 hours before or at least 4 hours after** you take COMPLERA.
- any of these medicines (if taken by mouth or injection):
 - clarithromycin (Biaxin)
 - erythromycin (E-Mycin, Eryc, Ery-Tab, PCE, Pediazole, Ilosone)
 - fluconazole (Diflucan)
 - itraconazole (Sporanox)
 - ketoconazole (Nizoral)
 - methadone (Dolophine)
 - posaconazole (Noxafil)
 - telithromycin (Ketek)
 - voriconazole (Vfend)
- medicines that are eliminated by the kidney, including acyclovir (Zovirax), cidofovir (Vistide), ganciclovir (Cytovene IV, Vitrasert), valacyclovir (Valtrex), and valganciclovir (Valcyte)

What are the possible side effects of COMPLERA?

COMPLERA can cause serious side effects, including:

- **See “What is the most important information I should know about COMPLERA?”**
- **New or worse kidney problems, including kidney failure,** can happen in some people who take COMPLERA. Your healthcare provider should do blood tests to check your kidneys before starting treatment with COMPLERA. If you have had kidney problems in the past or need to take another medicine that can cause kidney problems, your healthcare provider may need to do blood tests to check your kidneys during your treatment with COMPLERA.
- **Depression or mood changes.** Tell your healthcare provider right away if you have any of the following symptoms:
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself

- **Change in liver enzymes.** People with a history of hepatitis B or C virus infection or who have certain liver enzyme changes may have an increased risk of developing new or worsening liver problems during treatment with COMPLERA. Liver problems can also happen during treatment with COMPLERA in people without a history of liver disease. Your healthcare provider may need to do tests to check your liver enzymes before and during treatment with COMPLERA.
- **Bone problems** can happen in some people who take COMPLERA. Bone problems include bone pain, softening or thinning (which may lead to fractures). Your healthcare provider may need to do additional tests to check your bones.
- **Changes in body fat** can happen in people taking HIV medicine. These changes may include increased amount of fat in the upper back and neck (“buffalo hump”), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms and face may also happen. The cause and long term health effect of these conditions are not known.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting your HIV medicine.

The most common side effects of COMPLERA include:

- trouble sleeping (insomnia)
- abnormal dreams
- headache
- dizziness
- diarrhea
- nausea
- rash
- tiredness
- depression

Additional common side effects include:

- vomiting
- stomach pain or discomfort
- skin discoloration (small spots or freckles)
- pain

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of COMPLERA. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 (1-800-332-1088).

How should I take COMPLERA?

- **Stay under the care of your healthcare provider during treatment with COMPLERA.**
- **Take COMPLERA exactly as your healthcare provider tells you to take it.**
- **Always take COMPLERA with a meal.** Taking COMPLERA with a meal is important to help get the right amount of medicine in your body. A protein drink does not replace a meal.
- Do not change your dose or stop taking COMPLERA without first talking with your healthcare provider. See your healthcare provider regularly while taking COMPLERA.
- If you miss a dose of COMPLERA within 12 hours of the time you usually take it, take your dose of COMPLERA **with a meal** as soon as possible. Then, take your next dose of COMPLERA at the regularly scheduled time. If you miss a dose of COMPLERA by more than 12 hours of the time you usually take it, wait and then take the next dose of COMPLERA at the regularly scheduled time.
- Do not take more than your prescribed dose to make up for a missed dose.

This Brief Summary summarizes the most important information about COMPLERA. If you would like more information, talk with your healthcare provider. You can also ask your healthcare provider or pharmacist for information about COMPLERA that is written for health professionals, or call 1-800-445-3235 or go to www.COMPLERA.com

Issued: January 2013



COMPLERA, the COMPLERA Logo, EMTRIVA, GILEAD, the GILEAD Logo, GSI, HEPSERA, STRIBILD, TRUVADA, VIREAD, and VISTIDE are trademarks of Gilead Sciences, Inc., or its related companies. ATRIPLA is a trademark of Bristol-Myers Squibb & Gilead Sciences, LLC. All other marks referenced herein are the property of their respective owners.

©2013 Gilead Sciences, Inc. All rights reserved. CPAC0027 04/13

I COME FROM THE LAND DOWN UNDER

by: **Brad Loekle**



It's 10PM on the day after Sydney Mardi Gras and a very attractive, very high man in an adult diaper (with no shoes on) just face dived into the middle of a crowd of tables in a sidewalk café. Welcome to Australia & G'day, Mate!

I've been in Australia for almost a week and, the week before, was in New Zealand. I've wanted to come here for a very long time and finally, thanks to my endless travels on big gay cruise ships, my ship has come in down under. It is nothing like what I'd thought it would be and so much more than I could have ever imagined. And before we go any further, let me say now... it's a 24 hour flight there, and yes, it's worth it.

New Zealand is breathtaking beautiful. And not just one kind of beautiful, every kind of beautiful. Each city or region I travelled to was completely different from the next. One town looks like rural Scotland and the town just half a day's trip away looks like the lake region of Italy. It is so beautiful that you almost become immune to it and long to see a fat mid-western person or a Honey Boo Boo. There is one drawback to the beauty of the people of New Zealand. Particularly the men. Particularly the young men. A large number of them walk around barefoot. And I don't mean in the grass. I mean in the city streets. I don't know which came first, the barefooted Hobbits or the barefooted footed New Zealand men who live in the land where Hobbits were filmed... but either way: it's fucking gross. Call me old fashioned, but when I see a man walking barefoot through the streets of a city, I just think of tetanus and lock jaw, not free love and bohemia.

The men of Australia, on the other hand, keep their shoes on. Though it's rarely more than a flip flop. In fact, most of them, tanned and toned and perpetually sun kissed; seem to all dress as though they're going to a pool party in a John Hughes movie. Not that, that's a bad thing, but it's definitely a "thing". Also, and this I give you full warning of my Key West brethren and sistern: it is damn near impossible to get drunk on hard liquor in a bar in Australia. The

import taxes on booze are extremely high (imagine a bottle of Grey Goose at the liquor store costing almost \$100) and because of this they measure every shot. And every miniscule shot of booze is \$10+. This is probably why "Foster's is Australian for Beer"... because clearly, vodka is Australian for bankruptcy.

But even if you can't get drunk on booze, you can get drunk off eye candy. The men of Australia, in fact all the people of Australia, are some of the unilaterally pretty people I've ever seen. I mean, pretty. I mean, Hugh Jackman is like their Precious (and I mean the black girl, not the ring).

Oh and we have a lot of misconceptions about their country that I would like to now help sort out. Also this way you don't look like a damned fool (like I did) when you go there or meet one of them. So here are the important bullet points:

Koalas aren't bears, and they get REALLY pissy about this. The Koalas don't get pissy, but the people around them do. Koalas also don't get high off of the leaves they eat. It's eucalyptus not Purple Haze. The water does indeed go backwards in the drain there. It's poles thing. Only about half of them have their foreskin, so it's a really 50/50 shot as to what you get in the trouser department. Burger King is called Hungry Jack's... it has something to do with them hating the British but I don't know what that has to do with an American burger joint. "Pashes" means getting busy with someone. A "Loon" is a tranny chaser. And most of the gay community down there acts like they just discovered ecstasy, because it's all they seem to do and all they seem to have available. Again, not a bad thing, but definitely a "thing".

All in all, I'd say Australia is one of my new favorite places on earth. And at this point I've seen a lot of it. It's a hell of a long way from just about everywhere, but well worth the journey. Stay tuned because next month I'll be going to Bangkok, Thailand, Vietnam & China. These places I'm a little unsure of. Though it will be interesting to be in countries where, for the first time in my life, I would be considered a top!


More to cum....

Brad Loekle is a comedian currently living in LA. He can be seen Thursday nights on TruTV's "World's Dumbest" at 9PM & tours extensively on Atlantis Events & RSVP gay cruises.

For more on Brad, go to: www.bradhag.com

YOUR GLBT GUIDE TO

Key West



Q MAGAZINE

**THE ONLY LGBT
MAGAZINE IN THE KEYS**

Not only is Q Magazine is read by thousands of people in Key West, it's read on-line by people all over the world!

.....
ADVERTISING RATES
.....

Quarter Page Ad:	\$100 / Month
Half Page Ad:	\$200 / Month
Full Page Ad:	\$300 / Month
Inside Cover (front or back)	\$400 / Month
Outside Back Cover	\$450 / Month
Centerfold (two pages)	\$500 / Month

.....
**DISCOUNTS ARE OFFERED FOR COMMITMENT
CONTRACTS.**
.....

6 Month Contract	10% off monthly rate
12 Month Contract	15% off monthly rate

.....
CONTACT US:
305.396.8227 or info@keywestgayrag.com

www.QKeyWest.com

COMMANDO



MILITARY EVENT • ARMED FORCES WEEKEND

FRIDAY 5/17

"BALLS TO THE WALL" PARTY NIGHT
ALL U CAN DRINK DRAFT BEER \$10

SATURDAY 5/18

OFFICER'S POOL PARTY AFTERNOON
BBQ, FREE SHOTS, GAMES & GIVE AWAYS

DOG TAG DATING NIGHT
AMATEUR STRIP CONTEST AT MIDNIGHT

SUNDAY 5/19

U-DANCE 6-10PM
JESSICA DEVERAUX VARIETY SHOW 8PM

GO COMMANDO!

**★ NO COVER FOR THOSE WEARING
CAMOUFLAGE OR MILITARY ATTIRE**

MUSIC & VIDEOS BY
VJ/DJ GARY NOLAN
VJ/DJ NEIL

SOLDIER STIMULATION BY
THE MEN OF BOURBON

724 DUVAL • KEY WEST

Bourbon St. Pub

www.BourbonStPub.com

MEMORIAL DAY WEEKEND



BEACH BALL

FRIDAY, MAY 24

Kick-Off Party
Naked Splash NIGHT

SATURDAY, MAY 25

Beach Ball Pool Party AFTERNOON

Foam Party
Amateur Strip Contest NIGHT

SUNDAY, MAY 26

U-Dance 6-10PM

Jessica Deveraux Variety Show 8PM

Naked Splash NIGHT

MONDAY, MAY 27

Lazy Days Pool Party AFTERNOON

724 DUVAL • KEY WEST

Bourbon St. Pub

www.BourbonStPub.com

Pool Party



SATURDAYS NOON-5PM

**CLOTHING OPTIONAL | MULTIPLE SUNDECKS
PLAY AREAS | FREE SHOTS | HAPPY HOUR PRICES
POOL VOLLEYBALL | GAMES & PRIZES**

graffitti
Key West

MUSIC BY DJ NEIL



724 DUVAL • KEY WEST

Bourbon St. Pub

www.BourbonStPub.com


724 DUVAL • KEY WEST

New Orleans House

www.NewOrleansHouseKW.com

GULF OF MEXICO

KEY WEST HARBOR



ATLANTIC OCEAN

Q MAGAZINE welcomes you to America's most inclusive island!

Key West's official philosophy is "ONE HUMAN FAMILY" which means we believe everyone is entitled to equal rights, dignity and respect. You'll see a lot of these stickers, and you can get your own free "One Human Family" stickers at the Gay & Lesbian Visitor Center (513 Truman), the County Library (Fleming @ Elizabeth), or by sending a SASE to ONE HUMAN FAMILY, P.O. 972, Key West, FL 33041 or visit OneHumanFamily.info

ALL PEOPLE ARE CREATED EQUAL MEMBERS OF
ONE HUMAN FAMILY

You are invited to visit our valued advertisers!

- 1 Centennial Bank, 701 Whitehead & 1229 Simonton
- 2 Bourbon Street Pub - 724 Duval
- 3 801 Bourbon Bar - 801 Duval
- 4 Bobby's Monkey Bar - 900 Simonton
- 5 The Little Red Store - 901 Caroline Street
- 6 Equator Resort - 818 Fleming
- 7 Gay Trolley Starting Point - Angela & Duval
- 8 Big Ruby's Guesthouse - 409 Appelrouth Lane
- 9 Island House Resort for Men - 1129 Fleming
- 10 Alexander's Guest House - 1118 Fleming Street
- 11 KW Gay & Lesbian Visitor Center - 513 Truman
- 12 Monroe County Public Library - Fleming & Margaret
- 13 Truman Book & Video - 922 Truman
- 14 Graffiti for Men - 721 Duval
- 15 Amigos - 425 Green
- 16 In Touch - 706A Duval
- 17 Doug Mayberry Real Estate - 1075 Duval

ZACHARY MOSES : MAN ABOUT TOWN?

Chewy (the anonymous author of the Q magazine food column) is leaving us, and someone needs to fill his big hairy shoes.



Chewy and I met last Monday to go over the details of the food column. I will be the new author, and he was handing me the reins. We did this, appropriately, over dinner. The evening's choice? Café Sole, the intimate French restaurant at the corner of Frances and Southard.

The restaurant was packed; I would recommend having reservations before going. The hostess sat us at my favorite table; inside at the back next to the wine cabinet. The table is next to a wall of mirrors, which I can use to watch the goings on in the restaurant. What did I see first? Our muscle-bound, eastern-European waiter. You should have seen his forearms. He told us the specials, and I lost the ability to speak. When he departed, Chewy and I looked at each other and said "he's mine!"

We had no clue what his orientation was, but decided to assume he was family. We commenced with the cheesy compliments and senseless flirtations... did I mention he wears glasses? I love a man in glasses!

For dinner Chewy had fish, and I went with the duck. Café Sole is known for their fish, but I really wasn't feeling fish. I like to find out how well a restaurant handles their non-specialties, and the duck was fantastic! It was cooked to a perfect






crispiness, and I got to watch the waiter crack pepper all over it. When he came out holding that 2-foot-long phallus of a grinder, I decided I wanted pepper on everything! I had the man crack pepper on my crème brulee and strawberries. You know what? I enjoyed every bit of that peppered crème brulee too. Ever tried peppered strawberries? Unbelievably good!

After dinner, we decided we needed something a bit harder than the wine we'd had with dinner, so we got on our bikes and rode to Truman and Duval for the perfect St-Germain Martini at the Orchid Bar. This is the best place in town for this particular martini. The bar is very small, so cute patrons spill into the hotel's pool area. Why do I like it so much? Besides good drinks, it looks like the bar from the future. The colors, surfaces, the décor...I feel like this bar should be tended by robots. Plus, talk about cool; you fall into the pool, then grab a towel and say "I simply must get out of these wet clothes and into a dry martini!" And there would be a bartending robot, just waiting to pour you some St. Germain!

What's a man about town? Urban Dictionary says he is a sophisticated, suave urban dweller; he knows which bars to frequent, where the fashionistas dine, and which shows to see... is this me? Follow my column each month, and you decide. Bon Voyage Chewy!

Read more about Zachary's gay travel adventures at www.heTravel.com





jackadams

graffitti™
Key West
Since 1987



**721 Duval St.
305-295-0003**

AMIGOS

tortilla bar • key west



HOME OF THE
SQUARE TACO[®]

MEXICAN FOOD

breakfast ★ lunch ★ dinner

305.292.2009



"Once you've ruined
your reputation,
you can live
quite freely."

★ Key West ★

**ON
SALE!**

**DELIVERY
TO ALL OF
KEY WEST!
8 am - 11 pm**



Scan this code

Buy a T-SHIRT, and
we'll deliver it, too!

and "Like" us on Facebook
to receive a special offer!

425 Greene Street, Key West, FL 33040 • www.amigoskeywest.com

**Gabana
Realty, Inc.**

302 Southard St. - Suite 102

*I am NOT just
another pretty face.*

*32 years of Sales &
Property Management*

Charles E. Lee
Broker

305-294-6259

charlesleecbs@aol.com

REALTOR

**Mabry
Binnicker**

(305) 304-5624



**Preferred
Properties**

Coastal Realty, Inc.

KwMabry@msn.com

Ask Nana



E-mail your questions to nana@keywestgayrag.com

Dear Nana,

There's a guy I really like... a friend of mine. We definitely have sexual chemistry & I really want us to take it to the next level and date. The only problem is that he has a boyfriend of many years. They're totally open (too open if you ask me) and my friend definitely plays the field a lot. I think it's because he's not happy in the relationship. I think if he was with someone he really wanted to be with... like me... then he might stop sleeping around so much. My problem is that I don't want to confront him about our feelings for each other and lose him as a friend. So what do you think I should do?

Wooooo Nelly! Let's pull back the reins on the crazy train for a second and think about some of the words coming out of your mouth. First of all, it is not your job to decide what relationships are working and which aren't. Particularly if one of your main criteria for that is how it impacts



your ability to get ham slammed by one of the men in the relationship. Next, you really don't know they're unhappy. Perhaps your friend is with this guy because he's the only one who allows him to have these flings, while trusting that your friend comes home to him at the end of the day? Maybe not... but you don't know. Also, you're all grown-ass people. If he wants you, or wants to leave his boyfriend; he will. Be very careful about the idea of "well, if he was with me, he'd be different". Leopards rarely change their spots... and cocks do so even less. A man who uses his dick as a divining rod will be hard pressed to change; even for true love. You need to stop watching "Love, Actually" every Saturday night and start reading the writing on the wall. And for God's sake, don't miss out on someone really good for you who might be right under your nose because you can't stop staring at your free-dicked friend.

Dear Nana,

There's this really hot guy who's insanely into me and I kinda don't know how to take it. I'm not ugly and I don't have low self esteem... but I also know that I'm nowhere near in the same kinda shape that he's in. He has that perfect gay guy body and an insanely big dick to match. And I'm, well, just a normal looking guy in every sense of the word. In fact, I'm a little doughy and I'm worried that he's a "chubby chaser". Am I over thinking this? Should I just go for it?

I'll tell you how to take it... up your ass, deary! I love that you're worried about ulterior motives, but if only we could all have the problem of being pursued by a hot, hung hunk of man?! You're also completely ignoring the fact that he might be attracted to you because he's attracted to, YOU. Stranger things have happened than a gay man acting like a grown up and falling in love with a whole person, not just a peck and a prick. And even if he happens to have a penchant for paunch, what difference does that make? You are clearly turned on by his chiseled bod, so why can't his particular turn on be a boy with a little meat on the bone? Also, not for nothing, but if he happens to be black or Latino this would not be an uncommon thing. Maybe it's cultural, but I've seen a lot of ethnic hotties on both sides of the fence who only seem to want fuck a fatty. So take it, bitch! Take it all!!



*Do you have problems? Of course you do!! Why not let Nana help you?! Send your questions to:
Nana@keywestgayrag.com.*

THE TRUMAN ADULT PRIVATE VIDEO ARCADE

"Where you don't have
to spend a wad to
spend a wad!"



922 TRUMAN AVENUE - KEY WEST, FL
(305) 295 - 0120
TRUMANADULT.COM

- BOOKS
- TOYS/LUBES
- DVDS
- WHIPPETS
- RUSH ☺

- SPACIOUS
PRIVATE
BOOTHS
FOR YOUR
DISCREET
PLEASURE



Not just Gay -

Gay!



New Tees

GIFTS - CARDS - & MORE



with you

706-A Duval Street



DOUG MAYBERRY
— REAL ESTATE —
AND PROPERTY MANAGEMENT

*We're Your
Community Realtors®
in Paradise!*

Doug Mayberry • Sabrina Acevedo
Dean Townsend • Keith Bland
Kent Ducote • Mary Thixton
Bobby Ciulla • Ginger King • Terry Huff
John Dell • Michael Philip

(305) 292-6155

1075 Duval, Duval Square, Suite C23

DougMayberry.com

SALES &
RENTALS

RESIDENTIAL
COMMERCIAL

SNAP



SNAP



SNAP



Photos By
Kenne'
Tucker





The InQuisitor

Spilling the hottest tea in Key West

Until our phone gets turned off for non payment, we'll be here to listen to the latest dish. We'd never use real names but I'll bet if you ask around, you'll find out who's doing who! If you have some hot gossip, email us at tea@qkeywest.com. We'll keep your name a secret because we know how to keep secrets!

We hear that there is a certain straight bartender that may in the process of switching teams. Apparently he's been swinging his bat in a coworkers infield.

Rumor has it that a certain bar's door person has done some porn. We haven't found the video yet but we will comb through hours of porn to find it!

Sounds like there is a new outdoor cruising area. We're not sure how the word got out but we've been told that you wouldn't want to light a blacklight in the area!

Seems that a certain local walked out of a store and "accidentally" forgot to pay for his items. He was very embarrassed that he forgot but we think he was actually embarrassed that he got caught. No charges were filed but now he has to shop(lift) elsewhere for his club clothes,

We watched someone walk into the men's room 4 times in one hour and twice with someone. You either have a very small bladder (and sometimes need help) or you're not as clean as you claim to be!

35,000

UNIQUE ITEMS!



901 Caroline St., Flagler Station,
Key West, FL **305.396.7582**

FREE VISOR with \$10 purchase or more
and this coupon



BBQ • BREWS & BLUES

**F*CK THE
CONSTRUCTION**

**WE DELIVER
FAST AND FREE!**

- MENTION THIS AD FOR
10% OFF
EXCLUDING ALCOHOL

2502 NORTH ROOSEVELT BLVD

WWW.KWBQQ.COM

305.295.7427

SNAP



SNAP




SNAP



Photos By
Larry
Blackburn



YOUR STATUS IS A SWAB AWAY.



OraQuick[®] HIV Testing

20 minute rapid test

It's fast...It's free...results in 20 minutes

No Needles, No Blood, Painfree...Oral Swab



Serving Key West
and the Florida Keys



Monday 12-5pm

Tuesday 11-2pm

Tuesday 5-6pm

Wednesday 5-7pm

Thursday 6:30-8pm

Mon - Fri 9am - 5pm

Trinity Presbyterian Church 717 Simonton St.

Metropolitan Community Church 1215 Petronia St.

Gordon Rollins Center - 1434 Kennedy Drive

MLK Pool 300 Catherine St.(2nd and 4th Wed.)

Elks Lodge 1107 Whitehead St.(2nd and 4th Thurs.)

Gordon Rollins Center - 1434 Kennedy Drive

To schedule an appointment at any location,
call 305-296-6196 or 305-293-3669



**We're just a
phone call away.**

*Anytime... about anything,
from a simple question to
emergency help.*

***(305) 296-HELP
or 211***

*Part of the National Suicide Prevention Lifeline
800-273-TALK (8255) Operating 24/7 for calls
In Spanish & for Veteran's Crisis Line*

Whore--Scopes

Aquarius

Jan 20 – Feb 18

The stars show a promising month for you. You will only shart twice this month and one of those will be at home so no one will know!

Pisces

Feb 19 – Mar 20

Old habits die hard but when shopping for clothes, don't look in the "small" section... Or the "medium".... Keep going..... Keep going.... Keep going....

Aries

Mar 21 – Apr 19

I'm not saying you're a bad housekeeper, but when the haz-mat team from Hoarders refuses to go in your home, maybe it's time to make some changes.

Taurus

Apr 20 – May 20

Maybe it's time to start thinking about your future, unless your goals are to accomplish nothing, in that case, you're right on track!

Gemini

May 21 – Jun 20

You seem to have a black cloud around you all the time. It's called your personality. Try to smile once in a while and actually interact with people.

Cancer

Jun 21 – Jul 22

If you spent as much time working as you do looking in a mirror, you'd be a millionaire by now!

Leo

Jul 23 – Aug 22

Don't worry about what other people are wearing. You can't afford it!

Virgo

Aug 23 – Sep 22

You'll be asked to be on the Maury show this month. No, it's clearly not for someone looking to find out who the father is but for trying to find out who gave him Gonorrhea. (It was you!)

Libra

Sep 23 – Oct 22

Unless you're planning on becoming a drag queen, it's time to cut back on the make-up. If you really want better color, try prying your fat ass off the sofa and get some sun!

Scorpio

Oct 23 – Nov 21

You deserve more out of life. Now get out and get a job and earn some money!

Sagittarius

Nov 22 – Dec 21

When you can't find the tree through the forest, it's time to manscape!

Capricorn

Dec 22 – Jan 19

If you ever plan on getting a husband, you may want to make some changes. Start with not drinking water before bed. Wetting the bed every night is not cute!

Bobby's monkey BAR



**Cheapest Gay
Drinks in
Town!**

**900 Simonton Street
305-294-2655**

Karaoke

**Mon, Thur, Fri, & Sun
Starting at 9:30 PM**

**FREE Wii
Video Games**

**Key West's
Favorite Local
Gay Hang Out**



ISLAND HOUSE

BEST GAY RESORT IN THE WORLD

RESTAURANT

BAR

POOL

GYM

SAUNA

STEAMROOM

MASSAGE

TWO JACUZZIS

SUNDECK

EROTIC VIDEO LOUNGE

LOCKERS

TOWELS

24 HOURS

MEN ONLY

ID REQUIRED



1129 Fleming • 305 294 6284 • islandhousekeywest.com

All of the facilities of Island House are open and available 24 hours every day. Men are welcome at the restaurant and bar any time. Daily resort passes are available for locals and men staying elsewhere. Inquire at the front desk.